



Weathering of glaciogene marine clays from West Greenland

Belmonte, Louise Josefine; Foged, Niels Nielsen; Ingeman-Nielsen, Thomas

Published in:
Proceedings of Euroclay 2015

Publication date:
2015

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):
Belmonte, L. J., Foged, N. N., & Ingeman-Nielsen, T. (2015). Weathering of glaciogene marine clays from West Greenland. In *Proceedings of Euroclay 2015*

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Weathering of glaciogene marine clays from West Greenland

Louise Josefine Belmonte¹

Niels Foged¹

Thomas Ingeman-Nielsen¹

¹ DTU Civil Engineering (Arctic Technology Centre), Brovej bygn. 118, 2800 Kgs. Lyngby, Denmark

(email: lojon@byg.dtu.dk)

Deposits of fine-grained marine glaciogene sediments have been observed throughout the previously glaciated parts of the Northern hemisphere. Their particles were formed during glacial erosion of bedrock and deposited in marine environments during the retreating phases of the last glaciation. Although the sediments are distributed over a large geographical area, their overall grain size distributions and qualitative mineralogies are similar. They are dominated by the < 60 µm fraction (clay and silt fractions) and contain only minor amount of the 60-2000 µm fraction (sand fraction) (Belmonte et al., in prep.; Gillott, 1979; Pederstad and Jørgensen, 1985). Their chemical compositions are dominated by SiO₂ (46-63 wt%), Al₂O₃ (14-23 wt%) and Fe₂O₃T (4-15 wt%) with an alkali-alkaline earth element content (i.e. the content of K₂O, Na₂O, MgO and CaO) of 12-16 wt % (Bentley and Smalley, 1978; Gillott, 1971; Locat et al., 1984; Pederstad and Jørgensen, 1985; Roaldset, 1972). They often consist of feldspars, quartz, amphibole, mica/illite, chlorite, expandable clay minerals (e.g. smectite, vermiculite and mixed layer clays) and their < 2 µm fraction (clay fraction) contain a high percentage of primary minerals such as quartz, feldspars and amphibole (Belmonte et al., in prep.; Foged, 1979; Bentley and Smalley, 1978; Gillott, 1971; Gillott, 1979; Locat, 1984; Pederstad and Jørgensen, 1985; Ramesh and d'Angeljan, 1995; Roaldset, 1972).

The presence of the expandable clay minerals raises the question of their origin, for which there are several possibilities. They could have resulted from pre-glacial or sub-glacial weathering of bedrock, weathering in the pro-glacial fluvial system, weathering under marine conditions or, for some, weathering after exposure to sub-aerial conditions.

In this study, marine sediments from West Greenland have been characterised according to their grain size, clay mineralogy, plastic properties and pore water chemistry. The samples were selected to represent environments above and below present day sea level, and with and without permafrost in order to investigate the influence of the different environments on the weathering properties. We argue that the content of expandable clay minerals in the clay fraction indicates that chemical weathering must have been an active process during glacial erosion of bedrock, in deposition in seawater and/or after subaerial exposure.

Belmonte, L.J., Foged, N.N. and Ingeman-Nielsen, T. (in prep) Characterisation and weathering properties of fine-grained marine sediments from West Greenland.

Bentley, S.P. and Smalley, I.J. (1978) Mineralogy of sensitive clays from Quebec. *Canadian Mineralogist*, 16, pp. 103-112.

Foged, N. (1979) Ingeniørgeologiske undersøgelser af kvartære marine leraflejringer på Vestgrønland, Licentiate (Ph.d.) dissertation, Technical University of Denmark, Lyngby, Denmark (in Danish).

Gillott, J.E. (1971) Mineralogy of Leda clay, *Canadian Mineralogist*, 10, pp. 767-811.

Gillott, J.E. (1979) Fabric, composition and properties of sensitive soils from Canada, Alaska and Norway. *Engineering geology*, 14, pp. 149-172.

Locat, J., Lefebvre, G. and Ballivy, G. (1984) Mineralogy, chemistry, and physical properties interrelationships of some sensitive clays from Eastern Canada. *Canadian Geotechnical Journal*, 21, pp. 530-540.

Pederstad, K. and Jørgensen, P. (1985) Weathering in a marine clay during postglacial time. *Clay minerals*, 20, pp. 477-491.

Ramesh, R., d'Anglejan, B. (1995) Mineralogy, Chemistry and particle size interrelationships in some Post-Glacial Marine Deposits of the St. Lawrence Lowlands. *Journal of Coastal Research*, 11 (4), pp. 1167-1179.

Roaldset, E. (1972) Mineralogy and geochemistry of Quaternary clays in the Numedal area, southern Norway, *Norsk Geologisk Tidsskrift* 52 (4), pp. 335-369.